arXiv:1503.05413v1 [math.GM] 13 Mar 2015

De- Moivre's and Euler Formulas for Matrices of Split Quaternions

Melek Erdoğdu *   Mustafa Özdemir

March 19, 2015

Abstract

In this paper, real matrix representations of split quaternions are examined in terms of the casual character of quaternion. Then, we give De-Moivre' s formula for real matrices of timelike and spacelike split quaternions, separately. Finally, we state the Euler theorem for real matrices of pure split quaternions.

Keywords: Split Quaternion, De-Moivre's Formula, Euler Formula.
MSC Classification: 30C35.

1 Introduction

Sir William Rowan Hamilton discovered the quaternions in 1843, which was one of the his best contribution made to mathematical science. This discovery is a way of extending complex numbers to higher dimensions. So the set of quaternions, which was introduced by Hamilton, can be represented as

$$\mathbb{H} = \{q = q_0 + q_1 i + q_2 j + q_3 k; \quad q_0, q_1, q_2, q_3 \in \mathbb{R}\}$$

where

$$i^2 = j^2 = k^2 = -1 \text{ and } ijk = -1.$$

The set of quaternions is a member of noncommutative division algebra [9].

We know that real quaternion algebra is isomorphic to a real 4×4 matrix algebra. The real matrix representations of quaternions are investigated in [15], [6] and [7]. Euler and De-Moivre formulas for real matrices associated with quaternions are studied in [8].

In 1849, James Cockle introduced the set of split quaternions, also known as coquaternions. The real algebra of split quaternions, denoted by $\widehat{\mathbb{H}}$, is a four dimensional vector space over the real field $\mathbb{R}$ of real numbers with a basis $\{1, i, j, k\}$ satisfying

$$i^2 = -1, \ j^2 = k^2 = 1 \text{ and } ijk = 1.$$

The set of split quaternions is noncommutative, too. Unlike quaternion algebra, the set of split quaternions contains zero divisor, nilpotent elements and nontrivial idempotents [10], [13].

Split quaternions is a recently developing topic. There are some studies related to geometric applications of split quaternions such as [10], [11], [14] and [12]. Particularly, the geometric and physical applications of quaternions require solving quaternionic equations. Therefore, there are many studies on quaternionic and split quaternionic equations. For example; the rearrangement method of solving two sided linear split quaternionic equations is given in [5] and De Moivre's formula is used to find the roots of split quaternion in work [13]. Furthermore, 2×2 complex matrix representation of any split quaternion is presented in [1] and the left and right real matrix representations of split quaternions are studied in [10].

In this paper, we will investigate real matrix representations of split quaternions. First, we present a brief introduction of split quaternions. Then, we examine real matrices associated with split quaternions depending on the casual character of the quaternion. Moreover, we give De-Moivre' s formula for real matrices of timelike and spacelike split quaternions, separately. Finally, we state the Euler theorem for real matrices of pure split quaternions.

---

*Corresponding Author

2 Preliminaries

In this section, we present an introduction to split quaternions for the necessary background [10], [13], [14].

2.1 Split quaternions

The set split quaternions can be represented as

$$\widehat{\mathbb{H}} = \{q = q_0 + q_1 i + q_2 j + q_3 k; \quad q_0, q_1, q_2, q_3 \in \mathbb{R}\}$$

where the product table is:

	1	i	j	k
1	1	i	j	k
i	i	-1	k	$-j$
j	j	$-k$	1	$-i$
k	k	j	i	1

We write any split quaternion in the form $q = q_0 + q_1 i + q_2 j + q_3 k = S_q + \overrightarrow{V_q}$ where $S_q = q_0$ denotes the scalar part of q and $\overrightarrow{V_q} = q_1 i + q_2 j + q_3 k$ denotes vector part of q. If $S_q = 0$ then q is called pure split quaternion. The set of pure split quaternions are identified with the Minkowski 3 space. The Minkowski 3 space is Euclidean 3 space with the Lorentzian inner product

$$\langle \overrightarrow{u}, \overrightarrow{v} \rangle_{\mathbb{L}} = -u_1 v_1 + u_2 v_2 + u_3 v_3$$

where $\overrightarrow{u} = (u_1, u_2, u_3)$ and $\overrightarrow{v} = (v_1, v_2, v_3) \in \mathbb{E}^3$ and denoted by $\mathbb{E}^3_1$. We say that a vector $\overrightarrow{u}$ in $\mathbb{E}^3_1$ is spacelike, timelike or null if $\langle \overrightarrow{u}, \overrightarrow{u} \rangle_{\mathbb{L}} > 0$, $\langle \overrightarrow{u}, \overrightarrow{u} \rangle_{\mathbb{L}} > 0$ or $\langle \overrightarrow{u}, \overrightarrow{u} \rangle_{\mathbb{L}} = 0$, respectively. The conjugate of a split quaternion $q = q_0 + q_1 i + q_2 j + q_3 k$ is denoted by $\overline{q}$ and it is $\overline{q} = S_q - \overrightarrow{V_q} = q_0 - q_1 i - q_3 j - q_4 k$. For any $p, q \in \widehat{\mathbb{H}}$, the sum and product of split quaternions p and q are

$$\begin{aligned} p + q &= S_p + S_q + \overrightarrow{V_p} + \overrightarrow{V_q}, \\ pq &= S_p S_q + \left\langle \overrightarrow{V_p}, \overrightarrow{V_q} \right\rangle_{\mathbb{L}} + S_p \overrightarrow{V_q} + S_q \overrightarrow{V_p} + \overrightarrow{V_p} \times_{\mathbb{L}} \overrightarrow{V_q}, \end{aligned}$$

respectively. Here $\times_{\mathbb{L}}$ denotes Lorentzian vector product and is defined as

$$\overrightarrow{u} \times_{\mathbb{L}} \overrightarrow{v} = \begin{vmatrix} -e_1 & e_2 & e_3 \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix},$$

for vectors $\overrightarrow{u} = (u_1, u_2, u_3)$ and $\overrightarrow{v} = (v_1, v_2, v_3)$ of Minkowski 3 space. And norm of the split quaternion q is defined by

$$N_q = \sqrt{|q\overline{q}|} = \sqrt{|q_0^2 + q_1^2 - q_2^2 - q_3^2|}.$$

If $N_q = 1$ then q is called unit split quaternion and $q_0 = q/N_q$ is a unit split quaternion for $N_q \neq 0$. And the product

$$I_q = q\overline{q} = \overline{q}q = q_0^2 + q_1^2 - q_2^2 - q_3^2$$

determines the character of a split quaternion. A split quaternion is spacelike, timelike or lightlike (null) if $I_q < 0$, $I_q > 0$ or $I_q = 0$, respectively. Polar forms of split quaternions are defined as follows:

i. Every timelike split quaternion with spacelike vector part can be written in the form

$$q = N_q(\cosh\theta + \overrightarrow{\varepsilon} \sinh\theta)$$

where $\overrightarrow{\varepsilon}$ is spacelike unit vector in $\mathbb{E}^3_1$.

ii. Every timelike split quaternion with timelike vector part can be written in the form

$$q = N_q(\cos\theta + \overrightarrow{\varepsilon} \sin\theta)$$

where $\overrightarrow{\varepsilon}$ is timelike unit vector in $\mathbb{E}^3_1$.

iii. Every spacelike split quaternion can be written in the form

$$q = N_q(\sinh\theta + \overrightarrow{\varepsilon} \cosh\theta)$$

where $\overrightarrow{\varepsilon}$ is spacelike unit vector in $\mathbb{E}^3_1$.

2.2 De-Moivre's formula for split quaternions

Euler and De-Moivre formula for complex numbers are generalized for quaternions in [2], [3]. Similarly to the discussions in [2] and [3], De-Moivre formula and roots of split quaternions are investigated in [13] by considering the casual characters of the quaternion. De-Moivre formulas for split quaternions, which are stated in [13], can be given as follows;

i. If $q = N_q(\cosh\theta + \overrightarrow{\varepsilon}\sinh\theta)$ be a timelike split quaternion with spacelike vector part then

$$q^n = (N_q)^n(\cosh n\theta + \overrightarrow{\varepsilon}\sinh n\theta),$$

ii. If $q = N_q(\cos\theta + \overrightarrow{\varepsilon}\sin\theta)$ be a timelike split quaternion with timelike vector part then

$$q^n = (N_q)^n(\cos n\theta + \overrightarrow{\varepsilon}\sin n\theta),$$

iii. If $q = N_q(\sinh\theta + \overrightarrow{\varepsilon}\cosh\theta)$ be a spacelike split quaternion then

$$q^n = \left\{ \begin{array}{ll} (N_q)^n(\sinh\theta + \overrightarrow{\varepsilon}\cosh\theta), & n \text{ is odd} \\ \\ (N_q)^n(\cosh n\theta + \overrightarrow{\varepsilon}\sinh n\theta) & n \text{ is even} \end{array} \right\}$$

for $n \in \mathbb{N}$.

2.3 4×4 Real matrix representations of split quaternions

For any $q \in \widehat{\mathbb{H}}$, consider the linear map

$$g_q : \widehat{\mathbb{H}} \to \widehat{\mathbb{H}}$$

defined as

$$g_q(p) = qp.$$

This map is bijective and for $q = q_0 + q_1 i + q_2 j + q_3 k \in \widehat{\mathbb{H}}$, we have

$$\begin{aligned} g_q(1) &= q_0 + q_1 i + q_2 j + q_3 k, \\ g_q(i) &= -q_1 + q_0 i + q_3 j - q_2 k, \\ g_q(j) &= q_2 + q_3 i + q_0 j + q_1 k, \\ g_q(k) &= q_3 - q_2 i - q_1 j + q_0 k. \end{aligned}$$

Using this map, we may define an isomorphism between $\widehat{\mathbb{H}}$ and algebra of the matrices

$$\left\{ \begin{bmatrix} q_0 & -q_1 & q_2 & q_3 \\ q_1 & q_0 & q_3 & -q_2 \\ q_2 & q_3 & q_0 & -q_1 \\ q_3 & -q_2 & q_1 & q_0 \end{bmatrix} : q_0, q_1, q_2, q_3 \in \mathbb{R} \right\}.$$

The above obtained 4×4 real matrix for the split quaternion q is denoted by

$$\mathbf{L}_q = \begin{bmatrix} q_0 & -q_1 & q_2 & q_3 \\ q_1 & q_0 & q_3 & -q_2 \\ q_2 & q_3 & q_0 & -q_1 \\ q_3 & -q_2 & q_1 & q_0 \end{bmatrix}$$

and called the left matrix representation of q.

Proposition 1 *[10] For any $p, q \in \widehat{\mathbb{H}}$ and $r \in \mathbb{R}$, the followings are satisfied;*

i) $\mathbf{L}_{p+q} = \mathbf{L}_p + \mathbf{L}_q$,

ii) $\mathbf{L}_{pq} = \mathbf{L}_p\mathbf{L}_q$,

iii) $\mathbf{L}_{rp} = r\mathbf{L}_p$,

iv) $\mathbf{L}_1 = \mathbf{I}_4$.

Similar to the previous discussion, consider the linear map

$$f_q : \widehat{\mathbb{H}} \to \widehat{\mathbb{H}}$$

defined as

$$f_q(p) = pq.$$

This map is also bijective and for $q = q_0 + q_1 i + q_2 j + q_3 k \in \widehat{\mathbb{H}}$, we have

$$\begin{aligned} f_q(1) &= q_0 + q_1 i + q_2 j + q_3 k, \\ f_q(i) &= -q_1 + q_0 i - q_3 j + q_2 k, \\ f_q(j) &= q_2 - q_3 i + q_0 j - q_1 k, \\ f_q(k) &= q_3 + q_2 i + q_1 j + q_0 k. \end{aligned}$$

By this map, we may define an isomorphism between $\widehat{\mathbb{H}}$ and algebra of the matrices

$$\left\{ \begin{bmatrix} q_0 & -q_1 & q_2 & q_3 \\ q_1 & q_0 & -q_3 & q_2 \\ q_2 & -q_3 & q_0 & q_1 \\ q_3 & q_2 & -q_1 & q_0 \end{bmatrix} : q_0, q_1, q_2, q_3 \in \mathbb{R} \right\}.$$

We denote above corresponding 4×4 real matrix for any split quaternion q by

$$\mathbf{R}_q = \begin{bmatrix} q_0 & -q_1 & q_2 & q_3 \\ q_1 & q_0 & -q_3 & q_2 \\ q_2 & -q_3 & q_0 & q_1 \\ q_3 & q_2 & -q_1 & q_0 \end{bmatrix}$$

and it is called the right matrix representation of q.

Proposition 2 *[10] For any $p, q \in \widehat{\mathbb{H}}$ and $r \in \mathbb{R}$, the followings are satisfied;*

i) $\mathbf{R}_{p+q} = \mathbf{R}_p + \mathbf{R}_q$,

ii) $\mathbf{R}_{pq} = \mathbf{R}_q \mathbf{R}_p$,

iii) $\mathbf{R}_{rp} = r\mathbf{R}_p$,

iv) $\mathbf{R}_1 = \mathbf{I}_4$.

3 De Moivre's formula for real matrices of split quaternions

Depending on the casual character of split quaternion, we can express real matrix representations of quaternion as follows:
i. Every timelike split quaternion $q. = N_q(\cosh\theta + \overrightarrow{\varepsilon}\sinh\theta)$ with spacelike vector part, we may write

$$\mathbf{L}_q = N_q \begin{bmatrix} \cosh\theta & -u_1\sinh\theta & u_2\sinh\theta & u_3\sinh\theta \\ u_1\sinh\theta & \cosh\theta & u_3\sinh\theta & -u_2\sinh\theta \\ u_2\sinh\theta & u_3\sinh\theta & \cosh\theta & -u_1\sinh\theta \\ u_3\sinh\theta & -u_2\sinh\theta & u_1\sinh\theta & \cosh\theta \end{bmatrix} = N_q[\cosh\theta\, \mathbf{I}_4 + \sinh\theta\, \mathbf{L}_{\overrightarrow{\varepsilon}}],$$

$$\mathbf{R}_q = N_q \begin{bmatrix} \cosh\theta & -u_1\sinh\theta & u_2\sinh\theta & u_3\sinh\theta \\ u_1\sinh\theta & \cosh\theta & -u_3\sinh\theta & u_2\sinh\theta \\ u_2\sinh\theta & -u_3\sinh\theta & \cosh\theta & u_1\sinh\theta \\ u_3\sinh\theta & u_2\sinh\theta & -u_1\sinh\theta & \cosh\theta \end{bmatrix} = N_q[\cosh\theta\, \mathbf{I}_4 + \sinh\theta\, \mathbf{R}_{\overrightarrow{\varepsilon}}]$$

where $\overrightarrow{\varepsilon} = (u_1, u_2, u_3)$ is spacelike unit vector in $\mathbb{E}_1^3$.

ii. Every timelike split quaternion $q{=}N_q(\cos\theta + \overrightarrow{\varepsilon}\sin\theta)$ with timelike vector part, we may write

$$\mathbf{L}_q = N_q \begin{bmatrix} \cos\theta & -u_1\sin\theta & u_2\sin\theta & u_3\sin\theta \\ u_1\sin\theta & \cos\theta & u_3\sin\theta & -u_2\sin\theta \\ u_2\sin\theta & u_3\sin\theta & \cos\theta & -u_1\sin\theta \\ u_3\sin\theta & -u_2\sin\theta & u_1\sin\theta & \cos\theta \end{bmatrix} = N_q[\cos\theta\ \mathbf{I}_4 + \sin\theta\ \mathbf{L}_{\overrightarrow{\varepsilon}}],$$

$$\mathbf{R}_q = N_q \begin{bmatrix} \cos\theta & -u_1\sin\theta & u_2\sin\theta & u_3\sin\theta \\ u_1\sin\theta & \cos\theta & -u_3\sin\theta & u_2\sin\theta \\ u_2\sin\theta & -u_3\sin\theta & \cos\theta & u_1\sin\theta \\ u_3\sin\theta & u_2\sin\theta & -u_1\sin\theta & \cos\theta \end{bmatrix} = N_q[\cos\theta\ \mathbf{I}_4 + \sin\theta\ \mathbf{R}_{\overrightarrow{\varepsilon}}]$$

where $\overrightarrow{\varepsilon} = (u_1, u_2, u_3)$ is timelike unit vector in $\mathbb{E}^3_1$.

iii. Every spacelike split quaternion $q = N_q(\sinh\theta + \overrightarrow{\varepsilon}\cosh\theta)$, we may write

$$\mathbf{L}_q = N_q \begin{bmatrix} \sinh\theta & -u_1\cosh\theta & u_2\cosh\theta & u_3\cosh\theta \\ u_1\cosh\theta & \sinh\theta & u_3\cosh\theta & -u_2\cosh\theta \\ u_2\cosh\theta & u_3\cosh\theta & \sinh\theta & -u_1\cosh\theta \\ u_3\cosh\theta & -u_2\cosh\theta & u_1\cosh\theta & \sinh\theta \end{bmatrix} = N_q[\sinh\theta\ \mathbf{I}_4 + \cosh\theta\ \mathbf{L}_{\overrightarrow{\varepsilon}}],$$

$$\mathbf{R}_q = N_q \begin{bmatrix} \sinh\theta & -u_1\cosh\theta & u_2\cosh\theta & u_3\cosh\theta \\ u_1\cosh\theta & \sinh\theta & -u_3\cosh\theta & u_2\cosh\theta \\ u_2\cosh\theta & -u_3\cosh\theta & \sinh\theta & u_1\cosh\theta \\ u_3\cosh\theta & u_2\cosh\theta & -u_1\cosh\theta & \sinh\theta \end{bmatrix} = N_q[\sinh\theta\ \mathbf{I}_4 + \cosh\theta\ \mathbf{R}_{\overrightarrow{\varepsilon}}]$$

where $\overrightarrow{\varepsilon} = (u_1, u_2, u_3)$ is spacelike unit vector in $\mathbb{E}^3_1$.

Theorem 3 *For any timelike split quaternion $q = N_q(\cosh\theta + \overrightarrow{\varepsilon}\sinh\theta)$ with spacelike vector part, we have*

$$[\mathbf{L}_q]^n = (N_q)^n \begin{bmatrix} \cosh n\theta & -u_1\sinh n\theta & u_2\sinh n\theta & u_3\sinh n\theta \\ u_1\sinh n\theta & \cosh n\theta & u_3\sinh n\theta & -u_2\sinh n\theta \\ u_2\sinh n\theta & u_3\sinh n\theta & \cosh n\theta & -u_1\sinh n\theta \\ u_3\sinh n\theta & -u_2\sinh n\theta & u_1\sinh n\theta & \cosh n\theta \end{bmatrix} = (N_q)^n[\cosh n\theta\ \mathbf{I}_4 + \sinh n\theta\ \mathbf{L}_{\overrightarrow{\varepsilon}}],$$

$$[\mathbf{R}_q]^n = (N_q)^n \begin{bmatrix} \cosh n\theta & -u_1\sinh n\theta & u_2\sinh n\theta & u_3\sinh n\theta \\ u_1\sinh n\theta & \cosh n\theta & -u_3\sinh n\theta & u_2\sinh n\theta \\ u_2\sinh n\theta & -u_3\sinh n\theta & \cosh n\theta & u_1\sinh n\theta \\ u_3\sinh n\theta & u_2\sinh n\theta & -u_1\sinh n\theta & \cosh n\theta \end{bmatrix} = (N_q)^n[\cosh n\theta\ \mathbf{I}_4 + \sinh n\theta\ \mathbf{R}_{\overrightarrow{\varepsilon}}]$$

where $\overrightarrow{\varepsilon} = (u_1, u_2, u_3)$ is spacelike unit vector in $\mathbb{E}^3_1$.

Proof. We use induction on positive integer n. Let $q{=}N_q(\cosh\theta + \overrightarrow{\varepsilon}\sinh\theta)$ be any timelike split quaternion where $\overrightarrow{\varepsilon} = (u_1, u_2, u_3)$ is spacelike unit vector in $\mathbb{E}^3_1$. Assume that

$$[\mathbf{L}_q]^n = (N_q)^n \begin{bmatrix} \cosh n\theta & -u_1\sinh n\theta & u_2\sinh n\theta & u_3\sinh n\theta \\ u_1\sinh n\theta & \cosh n\theta & u_3\sinh n\theta & -u_2\sinh n\theta \\ u_2\sinh n\theta & u_3\sinh n\theta & \cosh n\theta & -u_1\sinh n\theta \\ u_3\sinh n\theta & -u_2\sinh n\theta & u_1\sinh n\theta & \cosh n\theta \end{bmatrix}$$

$$= (N_q)^n[\cosh n\theta\ \mathbf{I}_4 + \sinh n\theta\ \mathbf{L}_{\overrightarrow{\varepsilon}}],$$

holds. Then

$$(\mathbf{L}_q)^{n+1} {=} (\mathbf{L}_q)^n\ \mathbf{L}_q$$

$$= {N_q}^{n+1} \begin{bmatrix} \cosh n\theta & -u_1\sinh n\theta & u_2\sinh n\theta & u_3\sinh n\theta \\ u_1\sinh n\theta & \cosh n\theta & u_3\sinh n\theta & -u_2\sinh n\theta \\ u_2\sinh n\theta & u_3\sinh n\theta & \cosh n\theta & -u_1\sinh n\theta \\ u_3\sinh n\theta & -u_2\sinh n\theta & u_1\sinh n\theta & \cosh n\theta \end{bmatrix} \begin{bmatrix} \cosh\theta & -u_1\sinh\theta & u_2\sinh\theta & u_3\sinh\theta \\ u_1\sinh\theta & \cosh\theta & u_3\sinh\theta & -u_2\sinh\theta \\ u_2\sinh\theta & u_3\sinh\theta & \cosh\theta & -u_1\sinh\theta \\ u_3\sinh\theta & -u_2\sinh\theta & u_1\sinh\theta & \cosh\theta \end{bmatrix}$$

Using the equality

$$-u_1^2 + u_2^2 + u_3^2 = 1,$$

and the identities

$$\cosh\theta \cosh n\theta + \sinh\theta \sinh n\theta = \cosh(n+1)\theta,$$

$$\cosh\theta \sinh n\theta + \sinh\theta \cosh n\theta = \sinh(n+1)\theta,$$

we get

$$(\mathbf{L}_q)^{n+1} = (N_q)^{n+1} \begin{bmatrix} \cosh(n+1)\theta & -u_1 \sinh(n+1)\theta & u_2 \sinh(n+1)\theta & u_3 \sinh(n+1)\theta \\ u_1 \sinh(n+1)\theta & \cosh(n+1)\theta & u_3 \sinh(n+1)\theta & -u_2 \sinh(n+1)\theta \\ u_2 \sinh(n+1)\theta & u_3 \sinh(n+1)\theta & \cosh(n+1)\theta & -u_1 \sinh(n+1)\theta \\ u_3 \sinh(n+1)\theta & -u_2 \sinh(n+1)\theta & u_1 \sinh(n+1)\theta & \cosh(n+1)\theta \end{bmatrix}$$

$$= (N_q)^{n+1} \left\{ \cosh(n+1)\theta \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} + \sinh(n+1)\theta \begin{bmatrix} 0 & -u_1 & u_2 & u_3 \\ u_1 & 0 & u_3 & -u_2 \\ u_2 & u_3 & 0 & -u_1 \\ u_3 & -u_2 & u_1 & 0 \end{bmatrix} \right\}$$

$$= (N_q)^{n+1} \left[\cosh(n+1)\theta \; \mathbf{I}_4 + \sinh(n+1)\theta \; \mathbf{L}_{\overrightarrow{\varepsilon}} \right].$$

Hence the first formula is true. Similarly suppose that

$$(\mathbf{R}_q)^n = (N_q)^n \begin{bmatrix} \cosh n\theta & -u_1 \sinh n\theta & u_2 \sinh n\theta & u_3 \sinh n\theta \\ u_1 \sinh n\theta & \cosh n\theta & -u_3 \sinh n\theta & u_2 \sinh n\theta \\ u_2 \sinh n\theta & -u_3 \sinh n\theta & \cosh n\theta & u_1 \sinh n\theta \\ u_3 \sinh n\theta & u_2 \sinh n\theta & -u_1 \sinh n\theta & \cosh n\theta \end{bmatrix}$$

$$= (N_q)^n [\cosh n\theta \; \mathbf{I}_4 + \sinh n\theta \; \mathbf{R}_{\overrightarrow{\varepsilon}}]$$

is true. Then

$$(\mathbf{R}_q)^{n+1} = (\mathbf{R}_q)^n \; \mathbf{R}_q$$

$$= N_q^{n+1} \begin{bmatrix} \cosh n\theta & -u_1 \sinh n\theta & u_2 \sinh n\theta & u_3 \sinh n\theta \\ u_1 \sinh n\theta & \cosh n\theta & -u_3 \sinh n\theta & u_2 \sinh n\theta \\ u_2 \sinh n\theta & -u_3 \sinh n\theta & \cosh n\theta & u_1 \sinh n\theta \\ u_3 \sinh n\theta & u_2 \sinh n\theta & -u_1 \sinh n\theta & \cosh n\theta \end{bmatrix} \begin{bmatrix} \cosh\theta & -u_1 \sinh\theta & u_2 \sinh\theta & u_3 \sinh\theta \\ u_1 \sinh\theta & \cosh\theta & -u_3 \sinh\theta & u_2 \sinh\theta \\ u_2 \sinh\theta & -u_3 \sinh\theta & \cosh\theta & u_1 \sinh\theta \\ u_3 \sinh\theta & u_2 \sinh\theta & -u_1 \sinh\theta & \cosh\theta \end{bmatrix}$$

Then we get

$$(\mathbf{R}_q)^{n+1} = (N_q)^{n+1} \begin{bmatrix} \cosh(n+1)\theta & -u_1 \sinh(n+1)\theta & u_2 \sinh(n+1)\theta & u_3 \sinh(n+1)\theta \\ u_1 \sinh(n+1)\theta & \cosh(n+1)\theta & -u_3 \sinh(n+1)\theta & u_2 \sinh(n+1)\theta \\ u_2 \sinh(n+1)\theta & -u_3 \sinh(n+1)\theta & \cosh(n+1)\theta & u_1 \sinh(n+1)\theta \\ u_3 \sinh(n+1)\theta & u_2 \sinh(n+1)\theta & -u_1 \sinh(n+1)\theta & \cosh(n+1)\theta \end{bmatrix}$$

$$= (N_q)^{n+1} \left\{ \cosh(n+1)\theta \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} + \sinh(n+1)\theta \begin{bmatrix} 0 & -u_1 & u_2 & u_3 \\ u_1 & 0 & -u_3 & u_2 \\ u_2 & -u_3 & 0 & u_1 \\ u_3 & u_2 & -u_1 & 0 \end{bmatrix} \right\}$$

$$= (N_q)^{n+1} \left[\cosh(n+1)\theta \; \mathbf{I}_4 + \sinh(n+1)\theta \; \mathbf{R}_{\overrightarrow{\varepsilon}} \right].$$

So, the second formula is also true. ■

Theorem 4 *For any timelike split quaternion* $q{=}N_q(\cos\theta{+}\overrightarrow{\varepsilon}\sin\theta)$ *with timelike vector part,*

$$(\mathbf{L}_q)^n{=}(N_q)^n\begin{bmatrix} \cos n\theta & -u_1\sin n\theta & u_2\sin n\theta & u_3\sin n\theta \\ u_1\sin n\theta & \cos n\theta & u_3\sin n\theta & -u_2\sin n\theta \\ u_2\sin n\theta & u_3\sin n\theta & \cos n\theta & -u_1\sin n\theta \\ u_3\sin n\theta & -u_2\sin n\theta & u_1\sin n\theta & \cos n\theta \end{bmatrix}{=}(N_q)^n[\cos n\theta\ \mathbf{I}_4{+}\sin n\theta\ \mathbf{L}_{\overrightarrow{\varepsilon}}],$$

$$(\mathbf{R}_q)^n{=}(N_q)^n\begin{bmatrix} \cos n\theta & -u_1\sin n\theta & u_2\sin n\theta & u_3\sin n\theta \\ u_1\sin n\theta & \cos n\theta & -u_3\sin n\theta & u_2\sin n\theta \\ u_2\sin n\theta & -u_3\sin n\theta & \cos n\theta & u_1\sin n\theta \\ u_3\sin n\theta & u_2\sin n\theta & -u_1\sin n\theta & \cos n\theta \end{bmatrix}{=}(N_q)^n[\cos n\theta\ \mathbf{I}_4{+}\sin n\theta\ \mathbf{R}_{\overrightarrow{\varepsilon}}]$$

where $\overrightarrow{\varepsilon}=(u_1,u_2,u_3)$ *is timelike unit vector in* $\mathbb{E}_1^3$.

Proof. We use induction on positive integer n. For any timelike split quaternion $q{=}N_q(\cos\theta{+}\overrightarrow{\varepsilon}\sin\theta)$ with timelike unit vector $\overrightarrow{\varepsilon}=(u_1,u_2,u_3)$ in $\mathbb{E}_1^3$. Suppose that

$$(\mathbf{L}_q)^n=(N_q)^n\begin{bmatrix} \cos n\theta & -u_1\sin n\theta & u_2\sin n\theta & u_3\sin n\theta \\ u_1\sin n\theta & \cos n\theta & u_3\sin n\theta & -u_2\sin n\theta \\ u_2\sin n\theta & u_3\sin n\theta & \cos n\theta & -u_1\sin n\theta \\ u_3\sin n\theta & -u_2\sin n\theta & u_1\sin n\theta & \cos n\theta \end{bmatrix}$$

$$=(N_q)^n[\cos n\theta\ \mathbf{I}_4{+}\sin n\theta\ \mathbf{L}_{\overrightarrow{\varepsilon}}],$$

holds. Then

$$(\mathbf{L}_q)^{n+1}=(\mathbf{L}_q)^n\ \mathbf{L}_q$$

$$=(N_q)^{n+1}\begin{bmatrix} \cos n\theta & -u_1\sin n\theta & u_2\sin n\theta & u_3\sin n\theta \\ u_1\sin n\theta & \cos n\theta & u_3\sin n\theta & -u_2\sin n\theta \\ u_2\sin n\theta & u_3\sin n\theta & \cos n\theta & -u_1\sin n\theta \\ u_3\sin n\theta & -u_2\sin n\theta & u_1\sin n\theta & \cos n\theta \end{bmatrix}\begin{bmatrix} \cos\theta & -u_1\sin\theta & u_2\sin\theta & u_3\sin\theta \\ u_1\sin\theta & \cos\theta & u_3\sin\theta & -u_2\sin\theta \\ u_2\sin\theta & u_3\sin\theta & \cos\theta & -u_1\sin\theta \\ u_3\sin\theta & -u_2\sin\theta & u_1\sin\theta & \cos\theta \end{bmatrix}$$

Using the equality

$$-u_1^2+u_2^2+u_3^2=-1,$$

and the identities

$$\cos\theta\cos n\theta-\sin\theta\sin n\theta=\cos(n+1)\theta,$$

$$\cos\theta\sin n\theta+\sin\theta\cos n\theta=\sin(n+1)\theta,$$

we get

$$(\mathbf{L}_q)^{n+1}=(N_q)^{n+1}\begin{bmatrix} \cos(n+1)\theta & -u_1\sin(n+1)\theta & u_2\sin(n+1)\theta & u_3\sin(n+1)\theta \\ u_1\sin(n+1)\theta & \cos(n+1)\theta & u_3\sin(n+1)\theta & -u_2\sin(n+1)\theta \\ u_2\sin(n+1)\theta & u_3\sin(n+1)\theta & \cos(n+1)\theta & -u_1\sin(n+1)\theta \\ u_3\sin(n+1)\theta & -u_2\sin(n+1)\theta & u_1\sin(n+1)\theta & \cos(n+1)\theta \end{bmatrix}$$

$$=(N_q)^{n+1}\left\{\cos(n+1)\theta\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}+\sin(n+1)\theta\begin{bmatrix} 0 & -u_1 & u_2 & u_3 \\ u_1 & 0 & u_3 & -u_2 \\ u_2 & u_3 & 0 & -u_1 \\ u_3 & -u_2 & u_1 & 0 \end{bmatrix}\right\}$$

$$=(N_q)^{n+1}\left[\cos(n+1)\theta\ \mathbf{I}_4+\sin(n+1)\theta\ \mathbf{L}_{\overrightarrow{\varepsilon}}\right].$$

Hence the first formula is true.

Assume that

$$(\mathbf{R}_q)^n = (N_q)^n \begin{bmatrix} \cos n\theta & -u_1 \sin n\theta & u_2 \sin n\theta & u_3 \sin n\theta \\ u_1 \sin n\theta & \cos n\theta & -u_3 \sin n\theta & u_2 \sin n\theta \\ u_2 \sin n\theta & -u_3 \sin n\theta & \cos n\theta & u_1 \sin n\theta \\ u_3 \sin n\theta & u_2 \sin n\theta & -u_1 \sin n\theta & \cos n\theta \end{bmatrix}$$

$$= (N_q)^n [\cos n\theta \; \mathbf{I}_4 + \sin n\theta \; \mathbf{R}_{\overrightarrow{\varepsilon}}]$$

is true. Then

$$(\mathbf{R}_q)^{n+1} = (\mathbf{R}_q)^n \; \mathbf{R}_q$$

$$= (N_q)^{n+1} \begin{bmatrix} \cos n\theta & -u_1 \sin n\theta & u_2 \sin n\theta & u_3 \sin n\theta \\ u_1 \sin n\theta & \cos n\theta & -u_3 \sin n\theta & u_2 \sin n\theta \\ u_2 \sin n\theta & -u_3 \sin n\theta & \cos n\theta & u_1 \sin n\theta \\ u_3 \sin n\theta & u_2 \sin n\theta & -u_1 \sin n\theta & \cos n\theta \end{bmatrix} \begin{bmatrix} \cos \theta & -u_1 \sin \theta & u_2 \sin \theta & u_3 \sin \theta \\ u_1 \sin \theta & \cos \theta & -u_3 \sin \theta & u_2 \sin \theta \\ u_2 \sin \theta & -u_3 \sin \theta & \cos \theta & u_1 \sin \theta \\ u_3 \sin \theta & u_2 \sin \theta & -u_1 \sin \theta & \cos \theta \end{bmatrix}$$

Then we get

$$(\mathbf{R}_q)^{n+1} = (N_q)^{n+1} \begin{bmatrix} \cos(n+1)\theta & -u_1 \sin(n+1)\theta & u_2 \sin(n+1)\theta & u_3 \sin(n+1)\theta \\ u_1 \sin(n+1)\theta & \cos(n+1)\theta & -u_3 \sin(n+1)\theta & u_2 \sin(n+1)\theta \\ u_2 \sin(n+1)\theta & -u_3 \sin(n+1)\theta & \cos(n+1)\theta & u_1 \sin(n+1)\theta \\ u_3 \sin(n+1)\theta & u_2 \sin(n+1)\theta & -u_1 \sin(n+1)\theta & \cos(n+1)\theta \end{bmatrix}$$

$$= (N_q)^{n+1} \left\{ \cos(n+1)\theta \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} + \sin(n+1)\theta \begin{bmatrix} 0 & -u_1 & u_2 & u_3 \\ u_1 & 0 & -u_3 & u_2 \\ u_2 & -u_3 & 0 & u_1 \\ u_3 & u_2 & -u_1 & 0 \end{bmatrix} \right\}$$

$$= (N_q)^{n+1} [\cos(n+1)\theta \; \mathbf{I}_4 + \sin(n+1)\theta \; \mathbf{R}_{\overrightarrow{\varepsilon}}] \, .$$

So, the second formula is also true. ■

Theorem 5 *For any spacelike split quaternion* $q = N_q(\sinh\theta + \overrightarrow{\varepsilon} \cosh\theta)$*,*
If n odd then

$$(\mathbf{L}_q)^n = (N_q)^n \begin{bmatrix} \sinh n\theta & -u_1 \cosh n\theta & u_2 \cosh n\theta & u_3 \cosh n\theta \\ u_1 \cosh n\theta & \sinh n\theta & u_3 \cosh n\theta & -u_2 \cosh n\theta \\ u_2 \cosh n\theta & u_3 \cosh n\theta & \sinh n\theta & -u_1 \cosh n\theta \\ u_3 \cosh n\theta & -u_2 \cosh n\theta & u_1 \cosh n\theta & \sinh n\theta \end{bmatrix} = (N_q)^n [\sinh n\theta \; \mathbf{I}_4 + \cosh n\theta \; \mathbf{L}_{\overrightarrow{\varepsilon}}],$$

$$(\mathbf{R}_q)^n = (N_q)^n \begin{bmatrix} \sinh n\theta & -u_1 \cosh n\theta & u_2 \cosh n\theta & u_3 \cosh n\theta \\ u_1 \cosh n\theta & \sinh n\theta & -u_3 \cosh n\theta & u_2 \cosh n\theta \\ u_2 \cosh n\theta & -u_3 \cosh n\theta & \sinh n\theta & u_1 \cosh n\theta \\ u_3 \cosh n\theta & u_2 \cosh n\theta & -u_1 \cosh n\theta & \sinh n\theta \end{bmatrix} = (N_q)^n [\sinh n\theta \; \mathbf{I}_4 + \cosh n\theta \; \mathbf{L}_{\overrightarrow{\varepsilon}}].$$

If n is even then

$$(\mathbf{L}_q)^n = (N_q)^n \begin{bmatrix} \cosh n\theta & -u_1 \sinh n\theta & u_2 \sinh n\theta & u_3 \sinh n\theta \\ u_1 \sinh n\theta & \cosh n\theta & u_3 \sinh n\theta & -u_2 \sinh n\theta \\ u_2 \sinh n\theta & u_3 \sinh n\theta & \cosh n\theta & -u_1 \sinh n\theta \\ u_3 \sinh n\theta & -u_2 \sinh n\theta & u_1 \sinh n\theta & \cosh n\theta \end{bmatrix} = (N_q)^n [\cosh n\theta \; \mathbf{I}_4 + \sinh n\theta \; \mathbf{L}_{\overrightarrow{\varepsilon}}],$$

$$(\mathbf{R}_q)^n = (N_q)^n \begin{bmatrix} \cosh n\theta & -u_1 \sinh n\theta & u_2 \sinh n\theta & u_3 \sinh n\theta \\ u_1 \sinh n\theta & \cosh n\theta & -u_3 \sinh n\theta & u_2 \sinh n\theta \\ u_2 \sinh n\theta & -u_3 \sinh n\theta & \cosh n\theta & u_1 \sinh n\theta \\ u_3 \sinh n\theta & u_2 \sinh n\theta & -u_1 \sinh n\theta & \cosh n\theta \end{bmatrix} = (N_q)^n [\cosh n\theta \; \mathbf{I}_4 + \sinh n\theta \; \mathbf{L}_{\overrightarrow{\varepsilon}}].$$

where $\overrightarrow{\varepsilon} = (u_1, u_2, u_3)$ *is spacelike unit vector in* $\mathbb{E}_1^3$*.*

4 Euler formula for real matrices of split quaternions

In this part, we will state Euler formula for real matrices of spacelike and timelike pure split quaternions, separately.

Theorem 6 *For any timelike pure unit split quaternion* $q = \overrightarrow{\varepsilon} = (u_1, u_2, u_3)$,

$$e^{\theta \mathbf{L}_q} = \cos\theta \mathbf{I}_4 + \sin\theta\ \mathbf{L}_q \ \textit{and}\ e^{\theta \mathbf{R}_q} = \cos\theta \mathbf{I}_4 + \sin\theta\ \mathbf{R}_q.$$

Proof. Let $q = \overrightarrow{\varepsilon} = (u_1, u_2, u_3)$ be any timelike pure unit split quaternion, we have

$$\mathbf{L}_q = \begin{bmatrix} 0 & -u_1 & u_2 & u_3 \\ u_1 & 0 & u_3 & -u_2 \\ u_2 & u_3 & 0 & -u_1 \\ u_3 & -u_2 & u_1 & 0 \end{bmatrix} \text{ and } (\mathbf{L}_q)^2 = (-u_1^2 + u_2^2 + u_3^2) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Using the identity

$$-u_1^2 + u_2^2 + u_3^2 = -1$$

we get $(\mathbf{L}_q)^2 = -\mathbf{I}_4$. So, we have

$$\begin{aligned} e^{\theta \mathbf{L}_q} &= \mathbf{I}_4 + \theta \mathbf{L}_q + \frac{\theta^2}{2!}(\mathbf{L}_q)^2 + \frac{\theta^3}{3!}(\mathbf{L}_q)^3 + \frac{\theta^4}{4!}(\mathbf{L}_q)^4 + \frac{\theta^5}{5!}(\mathbf{L}_q)^5 + \frac{\theta^6}{6!}(\mathbf{L}_q)^6 + \cdots \\ &= \mathbf{I}_4 + \theta \mathbf{L}_q + \frac{\theta^2}{2!}(-\mathbf{I}_4) + \frac{\theta^3}{3!}(-\mathbf{L}_q) + \frac{\theta^4}{4!}(\mathbf{I}_4) + \frac{\theta^5}{5!}(\mathbf{L}_q) + \frac{\theta^6}{6!}(-\mathbf{I}_4) + \cdots \\ &= (1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \frac{\theta^6}{6!} + \cdots)\mathbf{I}_4 + (\theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \cdots)\mathbf{L}_q \\ &= \cos\theta\ \mathbf{I}_4 + \sin\theta\ \mathbf{L}_q. \end{aligned}$$

It is easy to see $(\mathbf{R}_q)^2 = -\mathbf{I}_4$ and $e^{\theta \mathbf{R}_q} = \cos\theta \mathbf{I}_4 + \sin\theta\ \mathbf{R}_q$. ■

Theorem 7 *For any spacelike unit pure split quaternion* $q = \overrightarrow{\varepsilon} = (u_1, u_2, u_3)$,

$$e^{\theta \mathbf{L}_q} = \cosh\theta\ \mathbf{I}_4 + \sinh\theta\ \mathbf{L}_q \ \textit{and}\ e^{\theta \mathbf{R}_q} = \cosh\theta\ \mathbf{I}_4 + \sinh\theta\ \mathbf{R}_q$$

Proof. Let $q = \overrightarrow{\varepsilon} = (u_1, u_2, u_3)$ be any spacelike pure unit split quaternion, we have

$$(\mathbf{L}_q)^2 = (-u_1^2 + u_2^2 + u_3^2) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Using the identity

$$-u_1^2 + u_2^2 + u_3^2 = 1$$

we get $(\mathbf{L}_q)^2 = \mathbf{I}_4$. So, we have

$$\begin{aligned} e^{\theta \mathbf{L}_q} &= \mathbf{I}_4 + \theta \mathbf{L}_q + \frac{\theta^2}{2!}(\mathbf{L}_q)^2 + \frac{\theta^3}{3!}(\mathbf{L}_q)^3 + \frac{\theta^4}{4!}(\mathbf{L}_q)^4 + \frac{\theta^5}{5!}(\mathbf{L}_q)^5 + \frac{\theta^6}{6!}(\mathbf{L}_q)^6 + \cdots \\ &= \mathbf{I}_4 + \theta \mathbf{L}_q + \frac{\theta^2}{2!}\mathbf{I}_4 + \frac{\theta^3}{3!}\mathbf{L}_q + \frac{\theta^4}{4!}\mathbf{I}_4 + \frac{\theta^5}{5!}\mathbf{L}_q + \frac{\theta^6}{6!}\mathbf{I}_4 + \cdots \\ &= (1 + \frac{\theta^2}{2!} + \frac{\theta^4}{4!} + \frac{\theta^6}{6!} + \cdots)\mathbf{I}_4 + (\theta + \frac{\theta^3}{3!} + \frac{\theta^5}{5!} + \cdots)\mathbf{L}_q \\ &= \cosh\theta\ \mathbf{I}_4 + \sinh\theta\ \mathbf{L}_q. \end{aligned}$$

Similarly, it can be easily shown that $(\mathbf{R}_q)^2 = \mathbf{I}_4$ and $e^{\theta \mathbf{R}_q} = \cosh\theta \mathbf{I}_4 + \sinh\theta\ \mathbf{R}_q$. ■

References

[1] Y. Alagöz, K. H. Oral, S. Yüce, *Split Quaternion Matrices.* Miskolc Mathematical Notes **13** (2012), 223-232.

[2] E. Cho, *De Moivre's formula for quaternions.* Applied Mathematics Letters **11** (1998), 33–35.

[3] E. Cho, *Euler's formula and De Moivre's formula for quaternions.* Missouri Journal of Mathematical Sciences **11** (1999), 80–83.

[4] M. Erdoğdu, M. Özdemir, *On Eigenvalues of Split Quaternion Matrices.* Advances in Applied Clifford Algebras **23** (2013), 615-623.

[5] M. Erdoğdu, M. Özdemir, *Two sided Linear Split Quaternionic Equations.* Linear and Multilinear Algebra (2013) DOI:10.1080/03081087.2013.851196.

[6] R. W. Farebrother, J. Grob, S.-O. Troschke, *Matrix representation of quaternions.* Linear Algebra and its Applications **362** (2003), 251-255.

[7] J. Grob, G. Trenkler, S.-O. Troschke, *Quaternions: further contributions to a matrix oriented approach.* Linear Algebra and its Applications **326** (2001), 205-213.

[8] M. Jafari, H. Mortazaasl, Y. Yaylı, *De-Moivre's Formula for Matrices of Quaternions.* JP Journal of Algebra, Number Theory and Applications **21** (2011), 57-67.

[9] I. L. Kantor, A. S. Solodovnikov, Hypercomplex Numbers, An Elementary Introduction to Algebras, Springer-Verlag, 1989.

[10] L. Kula, Y. Yaylı, *Split Quaternions and Rotations in Semi Euclidean Space.* Journal of Korean Mathematical Society **44** (2007), 1313-1327.

[11] M. Özdemir, A.A. Ergin, *Some geometric applications of split quaternions.* Proc. 16th Int. Conf. Jangjeon Math. Soc. **16** (2005), 108-115.

[12] M. Özdemir, A.A. Ergin, *Rotations with unit timelike quaternions in Minkowski 3-space,* Journal of Geometry and Physics **56** (2006), 322-336.

[13] M. Özdemir, *The Roots of a Split Quaternion.* Applied Mathematics Letters **22** (2009), 258-263.

[14] M. Özdemir, M. Erdoğdu, H. Şimşek, *On the Eigenvalues and Eigenvectors of a Lorentzian Rotation Matrix by Using Split Quaternions.* Advances in Applied Clifford Algebras **24** (2014), 179-192.

[15] F. Zhang, *Quaternions and Matrices of Quaternions.* Linear Algebra and its Applications **251** (1997), 21-57.

Melek Erdoğdu
Department of Mathematics-Computer Sciences
Necmettin Erbakan University
42060 Konya, Turkey
e-mail: merdogdu@konya.edu.tr.

Mustafa Özdemir
Department of Mathematics
Akdeniz University
07058 Antalya, Turkey
e-mail: mozdemir@akdeniz.edu.tr.